

I. L. Shynkovenko<sup>1</sup>, T. V. Ilyina<sup>1</sup>, A. M. Kovalyova<sup>1</sup>, O. V. Goryacha<sup>1</sup>, O. I. Golembiovska<sup>2</sup>, O. M. Koshovyi<sup>1</sup>

<sup>1</sup>National University of Pharmacy

<sup>2</sup>Institute of Organic Chemistry of NAS of Ukraine

## Saponins of the extracts of *Galium aparine* and *Galium verum*

**Aim.** To determine the qualitative composition and the quantitative content of saponins in the liquid extracts of the herb of cleavers (*Galium aparine* L.) and lady's bedstraw (*Galium verum* L.).

**Materials and methods.** Extracts of the herb of cleavers and lady's bedstraw were obtained by the triple extraction of the raw material with 96 % ethanol when heating with the subsequent concentration of the combined extracts to the ratio of the raw material : the finished product of 1 : 1. Saponins were studied by the method of high-performance liquid chromatography (HPLC).

**Results and discussion.** As a result of the chromatographic study of extracts of the herb of cleavers and lady's bedstraw it has been found that they contain 6 and 7 saponins, respectively. The compounds belong to ursane (ursolic, euscaphic, tormentic acids and uvaol), oleanane (oleanolic acid) and lupane (betulin and lupeol) type. In the extract from the herb of cleavers the compounds of ursane derivatives (50 mg/ml) prevail, the dominant compound is euscaphic acid (3.34 mg/ml); in the extract from the herb of lady's bedstraw there are saponins of lupane derivatives (2.50 mg/ml), lupeol (1.60 mg/ml) is predominant.

**Conclusions.** The results obtained indicate the prospects for further in-depth study of the chemical composition and biological activity of the liquid extracts from the herb of cleavers and lady's bedstraw.

**Key words:** cleavers; lady's bedstraw; liquid extract; saponins; HPLC

І. Л. Шинковенко, Т. В. Ільїна, А. М. Ковальова, О. В. Горяча, О. І. Голембівська, О. М. Кошовий

### Сапоніни екстрактів *Galium aparine* та *Galium verum*

**Мета роботи** – одержання, дослідження якісного складу та вмісту сапонінів у екстрактах рідких трави підмаренника чіпкого (*Galium aparine* L.) та підмаренника справжнього (*Galium verum* L.).

**Матеріали та методи.** Екстракти трави підмаренника чіпкого та підмаренника справжнього отримували шляхом триразової екстракції сировини 96 % спиртом етиловим при нагріванні з подальшим концентруванням об'єднаних витяжок до співвідношення сировина : готовий продукт 1 : 1. Сапоніни досліджували методом вискоєфективної рідинної хроматографії (ВЕРХ).

**Результати та їх обговорення.** В результаті хроматографічного дослідження екстрактів трави підмаренника чіпкого та підмаренника справжнього виявлено, що в них містяться 6 та 7 сапонінів відповідно. Сполуки відносяться до урсанового (урсолова, еускафова, торментинова кислоти та уваол), олеананового (олеанолова кислота) та лупанового (бетулін та лупеол) типу. В екстракті з трави підмаренника чіпкого в цілому переважають сполуки урсанового ряду (50 мг/мл), домінуючою сполукою є еускафова кислота (3,34 мг/мл); у екстракті з трави підмаренника справжнього – сапоніни лупанового ряду (2,50 мг/мл), домінуючою сполукою є лупеол (1,60 мг/мл).

**Висновки.** Отримані результати свідчать про перспективність подальшого поглибленого дослідження хімічного складу та біологічної активності екстрактів рідких з трави підмаренника чіпкого та підмаренника справжнього.

**Ключові слова:** підмаренник чіпкий; підмаренник справжній; екстракт рідкий; сапоніни; ВЕРХ

І. Л. Шинковенко, Т. В. Ильина, А. М. Ковалева, О. В. Горячая, Е. И. Голембiovская, О. Н. Кошевой

### Сапонины экстрактов *Galium aparine* и *Galium verum*

**Цель работы** – получение, изучение качественного состава и содержания сапонинов экстрактов жидких травы подмаренника цепкого (*Galium aparine* L.) и подмаренника настоящего (*Galium verum* L.).

**Материалы и методы.** Экстракты травы подмаренника цепкого и подмаренника настоящего получали путем трехкратной экстракции сырья 96 % спиртом этиловым при нагревании с дальнейшим концентрированием объединенных извлечений до соотношения сырье : готовый продукт – 1 : 1. Сапонины исследовали методом высокоэффективной жидкостной хроматографии (ВЭЖХ).

**Результаты и их обсуждение.** При хроматографическом изучении экстрактов травы подмаренника цепкого и подмаренника настоящего выявлено, что в них содержатся 6 и 7 сапонинов соответственно. Соединения относятся к урсановому (урсоловая, эускафовая, торментиновая кислоты и уваол), олеанановому (олеаноловая кислота) и лупановому (бетулин и лупеол) типу. В экстракте из травы подмаренника цепкого больше содержится сапонинов типа урсана (3,50 мг/мл), а доминирующим соединением является эускафовая кислота (3,34 мг/мл). В экстракте из травы подмаренника настоящего преобладают сапонины типа лупана (2,50 мг/мл) и доминирующим соединением является лупеол (1,60 мг/мл).

**Выводы.** Полученные результаты свидетельствуют о перспективности дальнейшего углубленного изучения химического состава и фармакологической активности экстрактов жидких из травы подмаренника цепкого и подмаренника настоящего.

**Ключевые слова:** подмаренник цепкий; подмаренник настоящий; экстракт жидкий; сапонины; ВЭЖХ

Among the representatives of *Galium* L. genus of *Rubiaceae* Juss. family cleavers (*Galium aparine* L.) and lady's bedstraw (*Galium verum* L.) belong to plants with the largest raw material base, they are distributed throughout the temperate zone of Europe, Asia and North America [1].

The herb of cleavers and lady's bedstraw is used for liver diseases, edemas, involuntary urination, painful urination, kidney stone disease, inflammation of the kidneys and bladder, inflammation of the stomach, intestines. Oil infusions of the cleavers herb are used externally for skin diseases, rashes, eczema, shingles, skin cancer [2-6].

A considerable number of works of scientists in many countries is devoted to the study of the chemical composition of the herb of *G. aparine* and *G. verum* [7-10]. Mainly phenolic compounds and terpenoids in the composition of essential oils were studied. In the samples of the herb of *G. verum* from Moldova a steroid compound of  $\beta$ -sitosterol and triterpenoid – oleanolic acid were found [11], in the samples from China there were  $\alpha$ -amyrin derivatives: ursolic acid, ursolic aldehyde and rubifolic acid [12]. Among other types in the herb of *G. rivale* the derivatives of oleanolic acid – rivosides A-E and momordin IIb were identified [13-15].

Many scientific publications are devoted to the study of such pharmacological activity of triterpenic acids as anti-inflammatory, antimicrobial, antiviral, cytotoxic, as well as their effect on the cardiovascular system [16-20].

**The aim** of this work was to study saponins in the extracts of the herb of *G. aparine* and *G. verum*.

#### Materials and methods

The raw material was collected at the flowering stage of plants in June, 2017: the herb of cleavers – in the Botanical garden of the National University of Pharmacy (NUPh, Kharkiv, Ukraine), the herb of lady's bedstraw – near the village of Rus'ki Tyshky of the Kharkiv region.

To obtain the liquid extracts 10.0 g of the air-dry crushed raw material was placed in a flask, 100 ml of 96 % ethanol was added and heated with a reflux condenser on a boiling water bath for 30 min. After cooling the content of the flask was filtered, and the volume of the resulting solution was measured. The procedure was repeated twice under the same conditions. The filtrates were combined and concentrated under vacuum in a rotary vacuum evaporator to the ratio of the raw material – finished product of 1 : 1; thus, 10 ml of the liquid extract was obtained.

The qualitative composition and quantitative content of saponins were studied by the method of high-performance liquid chromatography (HPLC) on a Shimadzu LC20 Prominence liquid chromatograph in the modular system equipped with the 4-channel pump LC20AD, the column thermostat CTO20A, the automatic sampler SIL20A, the diode array detector SPD20A and Chem Station LC20 in the following chromatographic conditions: the X-Bridge C18 chromatographic column with the size of 150 × 4.6 mm and the particle size of 5  $\mu$ m (Waters company); the column temperature – 30 °C; the detection wavelength – 205 nm; the flow rate of the mobile phase – 1.0 ml/min; the injection volume – 20  $\mu$ l;

Table 1

The chromatographic analysis of solutions of the reference samples

| Compounds      | Retention time, min | $\lambda_{max}$ , nm |
|----------------|---------------------|----------------------|
| Euscaphic acid | 8.53                | 200                  |
| Tormentic acid | 12.68               | 200                  |
| Betulin        | 14.57               | 200, 234, 322        |
| Oleanolic acid | 16.25               | 200                  |
| Ursolic acid   | 17.29               | 200                  |
| Uvaol          | 22.80               | 200                  |
| Lupeol         | 48.13               | 203, 230             |

the mobile phase – methanol for HPLC : 0.2 % solution of ammonium acetate (pH 6.75) in the ratio of 80 : 20; the elution mode – isocratic.

Identification of the components was carried out by the retention time and compliance of UV spectra with reference substances [21-23]. Triterpenic saponins was detected at 205 nm based on their absorption maximum at 200-210 nm.

The quantitative determination of individual components in ethanol extracts of the raw material was carried out using the external solutions of reference samples (Tab. 1).

The content of substances in the liquid extracts was calculated by the formula:

$$X, \text{ mg/ml} = \frac{A_{pr} \cdot m_{st} \cdot P}{A_{st} \cdot V_{st} \cdot 100},$$

where:  $A_{pr}$  – is the peak area of the substance on the chromatogram of the test solution;  $A_{st}$  – is the peak area of the substance on the chromatogram of the reference solution;  $m_{st}$  – is the weight of the standard sample of the substance in the reference sample solution, mg;  $V_{st}$  – is dilution of the reference solution, ml;  $P$  – is purity of the reference sample, %.

#### Results and discussion

The liquid extracts obtained are a thick green liquid with a specific odor.

The chromatographic analysis of saponins in the extract from the herb of cleavers 20 saponins were found, 6 saponins of them were identified; in the extract from the herb of lady's bedstraw there were 26 saponins, 7 of them were identified (Fig. 1-2, Tab. 2).

The chromatographic separation of oleanolic acid and ursolic acid is often hindered by the similarity of their chemical structure. On this basis the optimal chromatographic conditions and the type of the stationary column phase were selected, they allowed to achieve the separation index of these pentacyclic triterpenic acids of approximately 1.2. The retention time for oleanolic acid was 16.247 min, for ursolic acid – 17.288 min (Fig. 3).

As a result of the studies conducted the content of 6 and 7 saponins, respectively, was identified and determined in the liquid extracts of the herb of cleavers and lady's bedstraw (Tab. 2).

The compounds identified belong to ursane (ursolic, euscaphic, tormentic acids and uvaol), oleanane




Fig. 1. The scheme of the chromatogram of saponins in the extract of Cleavers

Table 2

Saponins of the liquid extracts of the herb of cleavers and lady's bedstraw

| Compounds      | <i>Galium aparine</i> L. |  | <i>Galium verum</i> L. |  |
|----------------|--------------------------|--|------------------------|--|
|                | Retention time, min      | The content of the substance in the extract, mg/ml | Retention time, min    | The content of the substance in the extract, mg/ml |
| Euscaphic acid | 8.22                     | 3.34   | 8.44                   | 0.87   |
| Tormentic acid | 12.97                    | 0,02   | 13.03                  | 0.004  |
| Betulin        | 14.60                    | 0.62   | 14.45                  | 0.89   |
| Oleanolic acid | 16.30                    | 0,21   | 16.43                  | 0.48   |
| Ursolic acid   | 17.28                    | 0.13   | 17.47                  | 1.04   |
| Uvaol          | -                        | 0.00   | 22.98                  | 0.01   |
| Lupeol         | 49.68                    | 1.20   | 49.40                  | 1.60   |
| In total       |                          | 5.52   |                        | 4.894  |




Fig. 2. The scheme of the chromatogram of saponins in the extract of Lady's bedstraw




Fig. 3. The typical chromatogram of the mixture of reference compounds of ursolic acid and oleanolic acid for separation

(oleanolic acid) and lupane (betulin and lupeol) type. The UV spectra of compounds are presented in Tab. 3

It has been found that the qualitative composition of saponins in the liquid extracts of lady's bedstraw and cleavers is similar. Of saponins identified only the extract of lady's bedstraw contains uvaol. Quantitatively, eus-









caphic acid (3.34 mg/ml) prevails in the extract of cleavers, whereas in the extract of lady's bedstraw lupeol (1.60 mg/ml) is dominant.

In the extract of cleavers ursane derivatives are dominant; their content is 3.50 mg/ml, the content of lupane derivatives is 1.82 mg/ml, oleanane – 0.21 mg/ml.

Table 3

Characteristics of saponins identified in the extracts of the herb of cleavers and lady's bedstraw

| Compounds        | Two-dimensional structural formula | UV-spectrum |
|------------------|------------------------------------|-------------|
| 1                | 2                                  | 3           |
| Ursolic acid     |                                    |             |
| Euscaphic Acid   |                                    |             |
| Tormentonic acid |                                    |             |

| 1              | 2   | 3  |
|----------------|---|--|
| Uvaol          |    |    |
| Oleanolic acid |    |    |
| Betulin        |   |   |
| Lupeol         |  |  |

In the extract of lady's bedstraw lupane derivatives prevail; their content is 2.50 mg/ml, the content of ursane derivatives is 1.92 mg/ml, oleanane – 0.48 mg/ml.

The results of the study confirm the data of the previous works on the presence of triterpenoids of the ursane type – ursolic acid in lady's bedstraw [12].

For the first time euscaphic acid, tormentic acid, betulin and lupeol were found in cleavers and lady's bedstraw. Uvaol was identified for the first time in lady's bedstraw.

The presence of the dominant compounds – euscaphic acid, lupeol, ursolic acid and betulin in the extract of cleavers and lady's bedstraw creates the basis for predicting such types of the pharmacological activity as hypoglycemic, anti-inflammatory, antimicrobial, antiprotozoal, anti-tumor, antiviral, antiproliferative, anti-angiogenic and hypocholesterolemic ones [24-28].

The dominance of euscaphic acid in the liquid extract from the herb of cleavers suggests the expediency

of studying its hypoglycemic activity, while the dominance of lupeol together with ursolic acid in the liquid extract from the herb of lady's bedstraw indicates the prospects of the study of its anti-tumor action.

### CONCLUSIONS

1. In the liquid extracts obtained from the herb of cleavers and lady's bedstraw the content of 6 and 7 saponins, respectively, has been identified and determined; they belong to ursane (ursolic, euscaphic, tormentic acids and uvaol), oleanane (oleanolic acid) and lupane (betulin and lupeol) type. Uvaol has been detected only in the extract from the herb of lady's bedstraw.

2. In the extract of cleavers ursane derivatives are dominant; their content is 3.50 mg/ml. In the extract of lady's bedstraw lupane derivatives prevail; their content is 2.50 mg/ml.

**Conflict of Interests:** authors have no conflict of interests to declare.

## REFERENCES

1. The Plant List (2013). Version 1.1. Published on the Internet. – Available at : <http://www.theplantlist.org/tpl1.1/record/kew-85676>, <http://www.theplantlist.org/tpl1.1/record/kew-87863>
2. An ethnobotanical study on the usage of wild medicinal herbs from Kopaonik Mountain (Central Serbia) / S. Jarić, Z. Popović, M. Mačukanović-Jocić et al. // *J. Ethnopharmacol.* – 2007. – Vol. 111 (1). – P. 160–175. <https://doi.org/10.1016/j.jep.2006.11.007>
3. Ethnobotanical study on traditional uses of wild medicinal plants in Prokletije Mountains (Montenegro) / N. Menković, K. Šavikin, S. Tasić et al. // *J. Ethnopharmacol.* – 2011. – Vol. 133. – P. 97–107. <https://doi.org/10.1016/j.jep.2010.09.008>
4. Medical ethnobotany of the Albanian Alps in Kosovo / B. Mustafa, A. Hajdari, F. Krasniqi et al. // *J. Ethnobiol. Ethnomed.* – 2012. – Vol. 8. – P. 6. <https://doi.org/10.1186/1746-4269-8-6>
5. Tradicionalna upotreba, kemijski sastav i biološki učinci vrsta roda *Galium* L / M. Friščić, M. Š. Baglama, M. Milović et al. // *Farmaceutski glasnik: glasilo Hrvatskog farmaceutskog društva.* – 2018. – Vol. 74 (5). – P. 343–350.
6. Traditional medicinal plant knowledge among Albanians, Macedonians and Gorani in the Sharr Mountains (Republic of Macedonia) / B. Rexhepi, B. Mustafa, A. Hajdari et al. // *Genet. Resour. Crop Evol.* – 2013. – Vol. 60. – P. 2055–2080. <https://doi.org/10.1007/s10722-013-9974-3>
7. Al-Snafi, A. E. Chemical Constituents and Medical Importance of *Galium aparine* a Review / A. E. Al-Snafi // *IAJPS.* – 2018. – Vol. 5 (3). – P. 1739–1744. <https://doi.org/10.5281/zenodo.1210517>
8. Banthorpe, D. V. Novel anthraquinones from undifferentiated cell cultures of *Galium verum* / D. V. Banthorpe, J. J. White // *Phytochemistry.* – 1995. – Vol. 38. – P. 107–111. [https://doi.org/10.1016/0031-9422\(94\)00579-i](https://doi.org/10.1016/0031-9422(94)00579-i)
9. Comparative study of polyphenolic content, antioxidant and antimicrobial activity of four *Galium* species (Rubiaceae) / L. Vlase, A. Mocan, D. Hanganu et al. // *Dig. J. Nanomater. Biostruct.* – 2014. – Vol. 9. – P. 1085–1094.
10. Composition of the essential oils of *Galium aparine* L. and *Galium odoratum* [L.] Scop. from Turkey / K. H. C. Baser, T. Ozek, N. A. Kirimer, L. Ergun. // *J. of Essential Oil Res.* – 2004. – Vol. 16 (4). – P. 305–307. <https://doi.org/10.1080/10412905.2004.9698728>
11. Investigations regarding the phytochemical study of some samples of *Galium verum* L. and *Galium album* Mill / G. Ghiță, R. Necula, A. Trifan et al. // *An Stiint Univ. Al I Cuza Iasi sect II a Biol. veget.* – 2012. – Vol. 58. – P. 45–50.
12. Chemical constituents of *Galium verum* / C. Zhao, J. Shao, D. Cao et al. // *Zhongguo Zhong Yao Za Zhi.* – 2009. – Vol. 34 (21). – P. 2761–2764.
13. Triterpene saponins and iridoid glucosides from *Galium rivale* / S. De Rosa, C. Iodice, M. Mitova et al. // *Phytochem.* – 2000. – Vol. 54 (8). – P. 1751–1756. [https://doi.org/10.1016/S0031-9422\(00\)00149-7](https://doi.org/10.1016/S0031-9422(00)00149-7)
14. Rivalosides A and B, two 19-Oxo triterpenoid saponins from *Galium rivale* / S. De Rosa, M. Mitova, N. Handjieva et al. // *J. Nat. Prod.* – 2000. – Vol. 63 (7). – P. 1012–1014. <https://doi.org/10.1021/np000073w>
15. Iridoid patterns in *Galium* L. and some phylogenetic considerations / M. I. Mitova, M. E. Anchev, N. V. Handjieva, S. S. Popov // *Z. Naturforsch. C.* – 2002. – Vol. 57. – P. 226–234. <https://doi.org/10.1515/znc-2002-3-405>
16. Connolly, J. Triterpenoids / J. Connolly, R. Hill // *Nat. Prod. Rep.* – 2008. – Vol. 25. – P. 794–830. <https://doi.org/10.1039/c6np00094k>
17. Inhibition of Skin Tumorigenesis by Rosemary and Its Constituents Carnosol and Ursolic Acid / M. T. Huang, C. H. Ho, Z. Y. Wang et al. // *Cancer Res.* – 1994. – Vol. 54. – P. 701–708.
18. Vechia, L. D. Oleanane and ursane derivatives and their importance on the discovery of potential antitumour, antiinflammatory and antioxidant drugs / L. D. Vechia, S. C. B. Gnoatto, G. Gosmann // *Quim. Nova.* – 2009. – Vol. 32. – P. 1245–1252. <https://doi.org/10.1590/S0100-40422009000500031>
19. Oleanolic Acid, a Pentacyclic Triterpene Attenuates the Mustard Oil-Induced Colonic Nociception in Mice / J. L. Maia, R. C. P. Lima-Junior, J. P. David et al. // *Biol. Pharm. Bull.* – 2006. – Vol. 29. – P. 82–85. <https://doi.org/10.1248/bpb.29.8>
20. Oleanolic acid, a pentacyclic triterpene attenuates capsaicin-induced nociception in mice: Possible mechanisms / J. L. Maia, R. C. P. Lima-Junior, C. M. Melo et al. // *Pharmacol. Res.* – 2006. – Vol. 54. – P. 282–286. <https://doi.org/10.1016/j.phrs.2006.06.003>
21. Leng, G. Determination of oleanolic acid and ursolic acid in different parts of *Mesona chinensis* Benth by RP HPLC / G. Leng // *Chin. J. Spectrosc. Lab.* – 2011. – Vol. 28. – P. 2111–2114.
22. Cen, J. H. High-performance liquid chromatographic analysis of bioactive triterpenes in *Perilla frutescens* / J. H. Cen, R. X. Xia, R. X. Tan // *J. Pharm. Biomed. Anal.* – 2003. – Vol. 32. – P. 1175–1179. [https://doi.org/10.1016/S0731-7085\(03\)00160-2](https://doi.org/10.1016/S0731-7085(03)00160-2)
23. Pentacyclic Triterpene Distribution in Various Plants Rich Sources for a New Group of Multi-Potent Plant Extracts / S. Jäger, H. Trojan, Th. Kopp et al. // *Molecules.* – 2009. – Vol. 14 (6). – P. 2016–2031. <https://doi.org/10.3390/molecules14062016>
24. Euscaphic acid, a new hypoglycemic natural product from folium *Eriobotryae* / J. Chen, W. L. Li, J. L. Wu et al. // *Pharmazie.* – 2008. – Vol. 63 (10). – P. 765–767.
25. Lupeol as a magical drug / A. Wal, R. S. Srivastava, P. Wal et al. // *Pharm. and Biol. Evaluations.* – 2015. – Vol. 2 (5). – P. 142–151.
26. Saleem, M. Lupeol, A Novel Anti-inflammatory and Anti-cancer Dietary Triterpene / M. Saleem // *Cancer Lett.* – 2009. – Vol. 285 (2). – P. 109–115. <https://doi.org/10.1016/j.canlet.2009.04.033>
27. Patočka, J. Biologically active pentacyclic triterpenes and their current medicine signification / J. Patočka // *J. of Applied Biomed.* – 2003. – Vol. 1. – P. 7–12.
28. Берштейн, Л. М. Урсоловая кислота как противоопухолевое средство и активатор белка-онкосупрессора PTEN и бурого жира / Л. М. Берштейн // *Вопросы онкол.* – 2012. – Т. 58, № 6. – С. 744–747.

## REFERENCES

1. The Plant List (2013). *Version 1.1*. Available at : <http://www.theplantlist.org/tpl1.1/record/kew-85676>, <http://www.theplantlist.org/tpl1.1/record/kew-87863>
2. Jarić, S., Popović, Z., Mačukanović-Jocić, M., Djurdjević, L., Mijatović, M., Karadžić, B., ... Pavlović, P. (2007). An ethnobotanical study on the usage of wild medicinal herbs from Kopaonik Mountain (Central Serbia). *Journal of Ethnopharmacology*, 111 (1), 160–175. <https://doi.org/10.1016/j.jep.2006.11.007>
3. Menković, N., Šavikin, K., Tasić, S., Zdunić, G., Stešević, D., Milosavljević, S., & Vincek, D. (2011). Ethnobotanical study on traditional uses of wild medicinal plants in Prokletije Mountains (Montenegro). *Journal of Ethnopharmacology*, 133 (1), 97–107. <https://doi.org/10.1016/j.jep.2010.09.008>

4. Mustafa, B., Hajdari, A., Krasniqi, F., Hoxha, E., Ademi, H., Quave, C. L., & Pieroni, A. (2012). Medical ethnobotany of the Albanian Alps in Kosovo. *Journal of Ethnobiology and Ethnomedicine*, 8 (1), 6. <https://doi.org/10.1186/1746-4269-8-6>
5. Friščić, M., Baglama, M. Š., Milović, M., Hazler, P. K., Maleš, Ž. (2018). Tradicionalna upotreba, kemijski sastav i biološki učinci vrsta roda Galium L. *Farmaceutski glasnik: glasilo Hrvatskog farmaceutskog društva*, 74 (5), 343–350.
6. Rexhepi, B., Mustafa, B., Hajdari, A., Rushidi-Rexhepi, J., Quave, C. L., & Pieroni, A. (2013). Traditional medicinal plant knowledge among Albanians, Macedonians and Gorani in the Sharr Mountains (Republic of Macedonia). *Genetic Resources and Crop Evolution*, 60 (7), 2055–2080. <https://doi.org/10.1007/s10722-013-9974-3>
7. Al-Snafi, A. E. (2018). Chemical Constituents and Medical Importance of Galium aparine a Review. *Indo American Journal of Pharmaceutical Sciences*, 5 (3), 1739–1744. <https://doi.org/10.5281/zenodo.1210517>
8. Banthorpe, D. V., & White, J. J. (1995). Novel anthraquinones from undifferentiated cell cultures of galium verum. *Phytochemistry*, 38 (1), 107–111. [https://doi.org/10.1016/0031-9422\(94\)00579-i](https://doi.org/10.1016/0031-9422(94)00579-i)
9. Vlase, L., Mocan A., Hanganu, D., Benedec, D. (2014). Comparative study of polyphenolic content, antioxidant and antimicrobial activity of four Galium species (Rubiaceae). *Dig. J. Nanomater Biostruct*, 9, 1085–1094.
10. Baser, K. H. C., Özek, T., Kirner, N., Deliorman, D., & Ergun, F. (2004). Composition of the Essential Oils of Galium aparineL. and Galium odoratum (L.) Scop. from Turkey. *Journal of Essential Oil Research*, 16 (4), 305–307. <https://doi.org/10.1080/10412905.2004.9698728>
11. Ghiță, G., Necula, R., Trifan, A., Zamfirache, M. M., Stănescu, U. (2012). Investigations regarding the phytochemical study of some samples of Galium verum L. and Galium album Mill. *An Stiint Univ Al I Cuza Iasi sect II a Biol. veget*, 58, 45–50.
12. Zhao, C., Shao, J., D, Cao., Zhang, Y., Li, X. (2009). Chemical constituents of Galium verum. *Zhongguo Zhong Yao Za Zhi*, 34 (21), 2761–2764.
13. De Rosa, S., Iodice, C., Mitova, M., Handjieva, N., Popov, S., & Anchev, M. (2000). Triterpene saponins and iridoid glucosides from Galium rivale. *Phytochemistry*, 54 (8), 751–756. [https://doi.org/10.1016/s0031-9422\(00\)00149-7](https://doi.org/10.1016/s0031-9422(00)00149-7)
14. De Rosa, S., Mitova, M., Handjieva, N., Popov, S., & Anchev, M. (2000). Rivalosides A and B, Two 19-Oxo Triterpenoid Saponins from Galium rivale. *Journal of Natural Products*, 63 (7), 1012–1014. <https://doi.org/10.1021/np000073w>
15. Mitova, M. I., Anchev, M. E., Handjieva, N. V., & Popov, S. S. (2002). Iridoid Patterns In Galium L. And Some Phylogenetic Considerations. *Zeitschrift für Naturforschung C*, 57 (3-4), 226–234. <https://doi.org/10.1515/znc-2002-3-405>
16. Hill, R. A., & Connolly, J. D. (2017). Triterpenoids. *Natural Product Reports*, 34 (1), 90–122. <https://doi.org/10.1039/c6np00094k>
17. Huang, M.-T., Ho, C.-H., Wang, Z. Y., Ferraro, T., Lou, Y. R., ... Conney, A. H. (1994). Inhibition of Skin Tumorigenesis by Rosemary and Its Constituents Carnosol and Ursolic Acid. *Cancer Res.*, 54, 701–708.
18. Dalla Vecchia, L., Gnoatto, S. C. B., & Gosmann, G. (2009). Derivados oleananos e ursanos e sua importância na descoberta de novos fármacos com atividade antitumoral, anti-inflamatória e antioxidante. *Química Nova*, 32 (5), 1245–1252. <https://doi.org/10.1590/s0100-40422009000500031>
19. Maia, J. L., Lima-Júnior, R. C. P., David, J. P., David, J. M., Santos, F. A., & Rao, V. S. (2006). Oleanolic Acid, a Pentacyclic Triterpene Attenuates the Mustard Oil-Induced Colonic Nociception in Mice. *Biological & Pharmaceutical Bulletin*, 29 (1), 82–85. <https://doi.org/10.1248/bpb.29.82>
20. Maia, J. L., Lima-Júnior, R. C. P., Melo, C. M., David, J. P., David, J. M., Campos, A. R., ... Rao, V. S. N. (2006). Oleanolic acid, a pentacyclic triterpene attenuates capsaicin-induced nociception in mice: Possible mechanisms. *Pharmacological Research*, 54 (4), 282–286. <https://doi.org/10.1016/j.phrs.2006.06.003>
21. Leng, G. (2011). Determination of oleanolic acid and ursolic acid in different parts of Mesona Chinensis Benth by RP HPLC. *Chin. J. Spectrosc. Lab.*, 28, 2111–2114.
22. Chen, J. H., Xia, Z. H., & Tan, R. X. (2003). High-performance liquid chromatographic analysis of bioactive triterpenes in Perilla frutescens. *Journal of Pharmaceutical and Biomedical Analysis*, 32 (6), 1175–1179. [https://doi.org/10.1016/s0731-7085\(03\)00160-2](https://doi.org/10.1016/s0731-7085(03)00160-2)
23. Jäger, S., Trojan, H., Kopp, T., Laszczyk, M., & Scheffler, A. (2009). Pentacyclic Triterpene Distribution in Various Plants – Rich Sources for a New Group of Multi-Potent Plant Extracts. *Molecules*, 14 (6), 2016–2031. <https://doi.org/10.3390/molecules14062016>
24. Wu, Y.-x., Jian, T.-y., Li, H., Ren, B. R., Zhang, H. Q. (2008). Euscaphic acid, a new hypoglycemic natural product from folium Eriobotryae. *Pharmazie*, 63 (10), 765–767.
25. Wal, A., Srivastava, R. S., Wal, P., Rai, A., Sharma, S. (2015). *Pharmaceutical and Biological Evaluations*, 2 (5), 142–151.
26. Saleem, M. (2009). Lupeol, a novel anti-inflammatory and anti-cancer dietary triterpene. *Cancer Letters*, 285 (2), 109–115. <https://doi.org/10.1016/j.canlet.2009.04.033>
27. Patočka J. (2003). Biologically active pentacyclic triterpenes and their current medicine signification. *Journal of Applied Biomedicine*, 1, 7–12.
28. Bershtein, L. M. (2012). *IIIoprosy onkologii*, 58 (6), 744–747.

**Information on authors:**

Shynkovenko I. L., postgraduate student of the Pharmacognosy Department, National University of Pharmacy

Ilyina T. V., Doctor of Pharmacy (Dr. habil.), professor of the Pharmacognosy Department, National University of Pharmacy. E-mail: [ilyinatany86@gmail.com](mailto:ilyinatany86@gmail.com).

ORCID: <http://orcid.org/0000-0003-3728-9752>

Goryacha O. V., Candidate of Pharmacy (Ph.D.), teaching assistant of the Pharmacognosy Department, National University of Pharmacy.

ORCID: <https://orcid.org/0000-0001-9877-7392>

Kovalyova A. M., Doctor of Pharmacy (Dr. habil.), professor of the Pharmacognosy Department, National University of Pharmacy. E-mail: [allapharm@yahoo.com](mailto:allapharm@yahoo.com).

ORCID: <http://orcid.org/0000-0002-1758-1222>

Golembiowska O. I., Candidate of Pharmacy (Ph.D.), engineer of the Organic Sulfur Chemistry Department, Laboratory of Condensed Heterocyclic Systems,

Institute of Organic Chemistry, National Academy of Sciences of Ukraine. E-mail: [golembiki@yahoo.fr](mailto:golembiki@yahoo.fr). ORCID: <https://orcid.org/0000-0001-5531-5374>

Koshovyi O. M., Doctor of Pharmacy (Dr. habil.), head of the Pharmacognosy Department, National University of Pharmacy

**Інформація про авторів:**

Шинковенко І. Л., аспірант кафедри фармакогнозії, Національний фармацевтичний університет

Ільїна Т. В., д-р фарм. наук, професор кафедри фармакогнозії, Національний фармацевтичний університет. E-mail: [ilyinatany86@gmail.com](mailto:ilyinatany86@gmail.com).

ORCID: <http://orcid.org/0000-0003-3728-9752>

Горяча О. В., канд. фарм. наук, асистент кафедри фармакогнозії, Національний фармацевтичний університет. ORCID: <https://orcid.org/0000-0001-9877-7392>

Ковальова А. М., д-р фарм. наук, професор кафедри фармакогнозії, Національний фармацевтичний університет. E-mail: [allapharm@yahoo.com](mailto:allapharm@yahoo.com).

ORCID: <http://orcid.org/0000-0002-1758-1222>

Голембіовська О. І., канд. фарм. наук, інженер відділу хімії органічних сполук сірки, лабораторія конденсованих гетероциклічних систем,

Інститут органічної хімії Національної академії наук України. E-mail: [golembiki@yahoo.fr](mailto:golembiki@yahoo.fr). ORCID: <https://orcid.org/0000-0001-5531-5374>

Кошовий О. М., д-р фарм. наук, завідувач кафедри фармакогнозії, Національний фармацевтичний університет

**Інформація об авторах:**

Шинковенко И. Л., аспирант кафедры фармакогнозии, Национальный фармацевтический университет

Ильина Т. В., д-р фарм. наук, профессор кафедры фармакогнозии, Национальный фармацевтический университет. E-mail: [ilyinatany86@gmail.com](mailto:ilyinatany86@gmail.com).

ORCID: <http://orcid.org/0000-0003-3728-9752>

Горячая О. В., канд. фарм. наук, ассистент кафедры фармакогнозии, Национальный фармацевтический университет.

ORCID: <https://orcid.org/0000-0001-9877-7392>

Ковалева А. М., д-р фарм. наук, профессор кафедры фармакогнозии, Национальный фармацевтический университет. E-mail: [allapharm@yahoo.com](mailto:allapharm@yahoo.com).

ORCID: <http://orcid.org/0000-0002-1758-1222>

Голембиовская Е. И., канд. фарм. наук, инженер отдела химии органических соединений серы, лаборатория конденсированных гетероциклических систем,

Институт органической химии Национальной академии наук Украины. E-mail: [golembiki@yahoo.fr](mailto:golembiki@yahoo.fr). ORCID: <https://orcid.org/0000-0001-5531-5374>

Кошевой О. Н., д-р фарм. наук, заведующий кафедрой фармакогнозии, Национальный фармацевтический университет

*Надійшла до редакції 14.08.2018 р.*